

RECEIVED
CENTRAL FAX CENTER
OCT 25 2007IN THE CLAIMS

1. (original) A lotion for therapy of dermatitis characterized in that it contains extracts drawn from one, two or more plants selected from the group consisting of Lightyellow Sophora Root, Turmeric, Magnolia Bark, Moutan Bark, Isatis Leaf, and Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn.f.).

Claim 2. (canceled)

3. (currently amended) The lotion for therapy of dermatitis described in claim 1, further comprising an auxiliary agent, characterized in that said auxiliary agent includes an auxiliary extract drawn from a medicinal herb, accelerator for skin permeation, accelerator for keratinization, alcohol and water.

4. (original) The lotion for therapy of dermatitis described in claim 3 characterized in that said auxiliary extract drawn from a medicinal herb contains extracts drawn from one, two or more selected from the group of Amur Cork Tree, Angelicae Dahurikae Root, Lemon, Smartweed and Licorice.

5. (withdrawn) The lotion for therapy of dermatitis described in claim 3 characterized in that said accelerating agent for skin permeation contains one, two or more selected from the group of Cnidii Rhizoma, Japanese Angelicae Root and dimethylsulfoxide.

6. (withdrawn) The lotion for therapy of dermatitis described in claim 3 characterized in that said accelerating agent for keratinization contains one or two selected from the group of

salicylic acid and resorcinol.

7. (withdrawn) The lotion for therapy of dermatitis described in claim 3 characterized in that the volume ratios of the extracts drawn from plants : accelerating agent for skin permeation : accelerating agent for keratinization in said auxiliary agent is 53 to 89% : 8 to 38%: 6 to 10%.

Claim 8. (canceled)

9. (previously presented) A lotion for therapy of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root, 2.7 to 3.3%; Turmeric, 1.8 to 2.2%; Magnolia Bark, 1.8 to 3.3%; Moutan Bark, 1.8 to 3.3%; Isatis Leaf, 0.9 to 1.1%; Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn.f.), 0.9 to 1.1%; Amur Cork Tree, 1.8 to 2.2%; Angelicae Dahuricae Root, 0.9 to 1.1%; Lemon, 0 to 3.3%; Smartweed, 0 to 2.2%; Licorice, 0 to 1%; Cnidii Rhizoma, 0.45 to 0.55%; Japanese Angelica Root, 0 to 0.55%; salicylic acid, 0.45 to 0.55%; resorcinol, 0.45 to 0.55%; alcohol, 25 to 35%; and water, 45 to 51%.

10. (previously presented) A lotion for therapy of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root, 3%; Turmeric, 2%; Magnolia Bark, 2%; Moutan Bark, 2%; Isatis Leaf, 1%; Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn.f.), 1%; Amur Cork Tree, 2%; Angelicae Dahuricae Root, 1%; Lemon, 3%; Smartweed, 2%; Licorice, 0.5%; Cnidii Rhizoma, 0.5%; Japanese Angelica Root, 0.5%; salicylic acid, 0.5%; resorcinol, 0.5%; alcohol, 30%; and water, 48.5%.

11. (previously presented) A lotion for therapy of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root, 2.7 to 3.3%; Turmeric, 1.8

-3-

Application No. 10/511,017

to 2.2%; Magnolia Bark, 1.8 to 2.2%; Moutan Bark, 1.8 to 2.2%; Isatis Leaf, 0.9 to 1.1%; Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn.f.), 0.9 to 1.1%; Amur Cork Tree, 1.8 to 2.2%; Angelicae Dahuricae Root, 0.9 to 1.1%; Lemon, 0 to 3.3%; Smartweed, 0 to 2.2%; Licorice, 0 to 0.55%; dimethylsulfoxide, 4.5 to 5.5%; salicylic acid, 0.45 to 0.55%; resorcinol, 0.45 to 0.55%; alcohol, 21 to 31%; and water, 45 to 51%.

12. (previously presented A lotion for therapy of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root, 3%; Turmeric, 2%; Magnolia Bark, 2%; Moutan Bark, 2%; Isatis Leaf, 1%; Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn.f.), 1%; Amur Cork Tree, 2%; Angelicae Dahuricae Root, 1%; Lemon, 3%; Smartweed, 2%; Licorice, 0.5%; dimethylsulfoxide, 5%; salicylic acid, 0.5%; resorcinol, 0.5%; alcohol, 26%; and water, 48.5%.